

## Drug Store

(405) 321-5300 2330 Mckown Norman OK

1008 Dr Tim Smith  
Tom Smith 2/10/2009  
AT 9AM AND 5PM

MAY CAUSE DROWSINESS

Avoid ALCOHOL-Do Not  
Drive if drowsy

1008

Tom Smith  
233 S Street  
Main CityDr Tim Smith  
123 Fake Stl  
TYLENOL/CODEINE#4 60-300 TAB 10  
00045051560

\$ 5.91

\$ 7.21

0 2/10/2009 MPH

(405) 444-4444

OK 22222-4444

Oklahoma City OK 73132-  
RR 0

\$ 7.21

White  
McNEILTYLENOL/CODEINE#4 60-300 TAB 10  
EXP Date 08-10-09  
May Refill 0 Times MPH

Round

Smith, Tom

## CODEINE(60) + APAP(300)

## WHY IS IT PRESCRIBED?

To relieve moderate to severe pain.

## BEFORE USING THIS MEDICINE:

Tell your doctor if you are having any of the following conditions:

KIDNEY DISEASE; CHRONIC OBSTRUCTIVE PULMONARY DISEASE (COPD); ASTHMA; LIVER PROBLEMS.

## HOW MEDICINE IS ADMINISTERED?

Use this medicine exactly as directed on the label, unless instructed differently by your doctor

\* This medicine can be taken with or without food, REGARDLESS of meal times. If stomach upset occurs, take with food.

\* DO NOT EXCEED recommended dosage; excessive use may lead to adverse effects or loss of effectiveness. Do not adjust the dose unless instructed.

## WHAT TO DO IF YOU MISS A DOSE?

This medicine is usually used as needed. If you miss a dose administer when or if needed. No double doses.

## HOW THIS MEDICINE SHOULD BE STORED?

Keep in original closed container in a dark, cool, and dry place away from children. Discard unneeded medicine.

## WARNINGS/PRECAUTIONS:

\* Safety of use during pregnancy has not been established. This medicine SHOULD NOT BE USED DURING PREGNANCY unless benefit justifies potential risk to the fetus. CALL YOUR DOCTOR.

\* This medicine may cause drowsiness, dizziness, or blurred vision. DO NOT drive or operate machines until you know how you react to this medicine.

\* AVOID concurrent use of alcohol or other CNS depressants as it may intensify the drowsiness, dizziness, or lightheadedness effect of this medicine.

\* This medicine may cause dizziness, lightheadedness or fainting. Avoid sudden changes in posture. BE CAREFUL and slowly sit up or stand up.

\* Prolonged use of this medication may cause physical and psychological dependence, do not exceed recommended dose.

\* Strictly limit alcohol consumption. LIVER PROBLEMS may be more likely to occur if you drink alcoholic beverages regularly while taking this medicine.

Stop taking this medicine and notify your doctor AS SOON AS POSSIBLE if you experience:

Yellowing of eyes or skin; Jerkings of face, trunk, or extremities; Shortness of breath; Skin rash; Irregular heartbeat; Low blood pressure; Hot sensation and redness of face.

## Other Common Side Effects:

Sedation; Nausea; Constipation; Dizziness; Vomiting; Dry mouth; Sweating; Anemia; Abnormal blood cell count; Changes in urination frequency or amount; Decreased appetite; Blurred vision; Confusion; Headache; Lightheadedness; Feeling of general discomfort; Drowsiness.

If symptoms are mild but do not go away or are bothersome, check with your doctor. IF ANY OF THE ABOVE SIDE EFFECTS IS SEVERE, CALL YOUR DOCTOR IMMEDIATELY.

Drug Store  
2330 Mckown

Norman

(405) 321-5300

OK 1 2

1008  
Tom Smith233 S Street  
Main City

OK 22222-4444

Dr Tim Smith  
2/10/2009

(405) 444-4444

TYLENOL/CODEINE#4 60-300 TAB 10

\$ 7.21

\$ 7.21